

Tissue Culture and Micropropagation of Mongolian Licorice (*Glycyrrhiza uralensis* Fisch.)

Yu. Oyunbileg¹, Kh. Altanzul¹, Ts. Oyunsuren¹, D. Bayarlkhagva²

¹Laboratory of Molecular Biology, Institute of Biology, Mongolian Academy of Sciences, Ulaanbaatar 210351, Mongolia, e-mail: yungeree@yahoo.com

²Department of Genetics and Molecular Biology, Faculty of Biology, National University of Mongolia, Ulaanbaatar 210646, Mongolia

Abstract

Mongolian licorice (*Glycyrrhiza uralensis* Fisch.) is a pharmacologically important plant rich in flavonoids and saponins. For tissue culture, root and cotyledon explants from seedlings were used. Sterilized explants with one node were used for micropropagation. Half-strength Murasige-Skuge medium and Gamborg's B5 medium with different supplements were used for the induction of calluses and multiple shoots. Conditions for tissue culture and micropropagation of *G. uralensis* were determined.

Key words: callus, *Glycyrrhiza uralensis*, *in vitro*, micropropagation, Mongolia, nodal culture

Introduction

Currently, there is much international interest in increasing plant resources, plant productivity and the ability to synthesize specific compounds, especially various secondary metabolites useful for medicinal practices.

Licorice (*Glycyrrhiza uralensis* Fisch.), belonging to the family *Fabaceae*, is recognised as one of the most valuable and widely used oriental herbs. During the last few years, use of licorice has increased rapidly. However, due to human influence and natural desertification processes in Mongolia licorice resources may become exhausted in the near future.

In vitro culture and propagation are useful tools in the conservation of this important plant. Use of *in vitro* cultures of *G. uralensis* and *G. glabra* L. have been reported by a number of authors (Thengane *et al.*, 1998; Kovalenko & Kurchii, 1998; Kohjyouma *et al.*, 1995). The roots and isolated active components of these plants are widely used for treatment of viral infections, inflammatory diseases and prevention of different cancers (Arase *et al.*, 1997; Numazaki, 2003; Shiraki *et al.*, 2003).

The aim of our research is to determine methods of *in vitro* culture and micropropagation of Mongolian licorice, in order to increase the bioresources of *G. uralensis* in Mongolia.

Materials and Methods

Plant materials. Seed samples of *G. uralensis* were provided by researchers from the Institute of Botany at the Mongolian Academy of Sciences. Two and four year old *G. uralensis* leaves, roots and whole plants were also collected from Dashinchilen district in Bulgan province, and Bogd and Baatsagaan districts in Bayankhongor province during a field study.

Culture conditions. The seeds were soaked in sterilized water for 24 hours, followed by a 70% ethanol bath for 30 seconds. Stratifications were performed at 4°C for 72 hours, and the seed surface sterilized with 2% sodium hypochloride solution with a drop of Tween-20 for 5 minutes, before being rinsed with sterile water 3 times. The prepared seeds were then used for germination and micropropagation experiments.

For germination of *G. uralensis*, half-strength basal Murashige-Skoog medium (MS medium; pH 5.8) supplemented with 0.8 % agar and 1.5 % sucrose was used. Autoclaving was performed at +121 °C for 15 minutes (Oyunbileg & Mijidsuren, 2004).

Seedlings grown on Gamborg's B5 medium and supplemented with 4 different combinations of growth regulators (BAP-1.0 mg/L and 2.0 mg/L, IAA-0.1 mg/L and 1.0 mg/L) (Liao *et al.*, 2004) were used for micropropagation.

Incubation. All cultures were incubated at 25 °C under 8/16 hours photoperiod in a daylight cabinet. Light intensity was 40 $\mu\text{Mole}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$.

Results and Discussion

Naturally grown licorice has poor seed germination, about 1 % to 8 % (Gankhuyag & Ligaa, 1993; Dashjamts, 1983). Gankhuyag and Ligaa (1993) stated that it is possible to achieve 92 % seed germination through scarification, and previous researchers also determined that seed germination could be increased after mechanical damage to the seed coat (Gankhuyag & Ligaa, 1993; Dashjamts, 1983). In this study, seed coats were damaged and the stratification method was used, as a result 55.8 % - 100 % of seeds germinated. There was no significant difference between mediums (1/2 MS, 1/2 B5, 1 % agar) used in the experiment as shown in Figure 1.

One week after germination, root and cotyledon explants from seedlings were transferred to

with those initiated from cotyledon explants (Fig. 2).

Propagation of *G. uralensis* was performed using nodal and shoot tip cultures, on Gamborg's B5 medium, which was supplemented with 5 different combinations of BAP and IAA growth regulators. There was no significant difference between the lengths of shoots and the addition of BAP and IAA combinations did not increase shoot formation (Table 2).

In addition, the seedlings were grown on half-strength MS medium supplemented with 4 different combinations of BAP and NAA growth regulators. The results showed that half-strength MS medium with 0.1 mg/l NAA was effective in inducing the root of licorice (Table 3). After 14 days of culture, the medium with no growth regulators was effective in the formation of multiple shoots and roots. Some shoots rooted spontaneously on half-strength MS medium without growth regulators. This medium was therefore selected for rooting and regeneration of whole plantlets. As a result, after 42 days of




Figure 1. Comparative seed germination of *G. uralensis* (in 1/2 MS, 1/2 B5, 1 % agar mediums) after one week, collected from 3 different locations.

Gamborg's B5 medium and supplemented with 1 mg/l 2,4-dichlorophenoxyacetic acid (2,4-D), 0.1 mg/l kinetin and 2% sucrose to initiate callus growth. Two to three weeks later, callus initiation occurred (Table 1).

Calluses that grew from root explants were more friable, yellowish in colour and watery compared

culture the plantlets developed (Fig. 3), but only 50 plantlets were transplanted into pots. Currently we are continuing our research, and regenerated plantlets are being acclimatized.

The nodal cultivated on MS medium formed new shoots after 10 days. In order to identify the optimal medium for regeneration, cultures from

Table 1. Callus initiation from root and cotyledon explants, after 3 weeks.

Location of sample	Explant	Number of tested explants	Number of induced calluses
Bulgan:	Cotyledon	30	25
Dashinchilen	Root	30	27
Bayankhongor:	Cotyledon	30	26
Baatsagaan	Root	30	28
Bayankhongor: Bogd	Cotyledon	30	25
	Root	30	28

Table 2. Effect of BAP and IAA on root formation and shoot development

Treatments	Concentration of growth regulators (mg/l)		Explants tested (N)	Root formation rate (%)	Length of shoot (cm)
	BAP	IAA			
1	0	0	9	55.6	2.72 ± 0.18
2	2.0	0.1	9	0	2.50 ± 0.25
3	2.0	1.0	9	0	2.66 ± 0.15
4	1.0	0.1	9	0	2.33 ± 0.27
5	1.0	1.0	9	0	2.61 ± 0.24




Figure 2. Callus grown from the cotyledon (a.) and the root (b.) explants.


Figure 3. A. Transferred shoot tips of *G. uralensis* into new 1/2 MS medium. B. After 6 weeks.

Figure 4. New shoot formation by nodal cultures and influence of secreted phenolics.

seedlings and multiple shoots from nodal cultures were sub-cultured on half-strength MS medium. The results of our experiments showed that regenerated plantlets could be obtained after about 3 weeks (Fig. 4). These results were similar to those conducted by Thengane *et al.* (1998) and Kohjyouma *et al.* (1995) who developed *in vitro* plantlets of *Glycyrrhiza glabra*.

One problem raised during the *in vitro* micropropagation was the secretion of phenol-like substances, which inhibited the growth of tissue culture. The same phenomenon was observed by Kovalenko *et al.* (1998). In this case, the secretion of phenolics was regulated by abscisic acid, which acts as an antioxidant. In our experiment, it was found that frequently transferring the new shoots

Table 3. Effect of BAP and NAA on root formation and shoot development

Treatments	Concentration of growth regulators (mg/l)		Explants tested (N)	Root formation rate (%)	Length of shoot (cm)
	BAP	IAA			
1	0	0	9	55.6	2.72 ± 0.18
2	2.0	0.1	9	0	2.50 ± 0.25
3	2.0	1.0	9	0	2.66 ± 0.15
4	1.0	0.1	9	0	2.33 ± 0.27
5	1.0	1.0	9	0	2.61 ± 0.24

to new medium prevented the phenol-like substances from influencing culture growth.

The results obtained from this study would be useful for increasing *G. uralensis* resources and for secondary metabolism research.

Acknowledgements

We would like to thank the Asia Research Center at the National University of Mongolia and the Korea Foundation for Advanced Studies for supporting this research. Thanks also to Drs. D. Magsar and G. Ochirbat from the Institute of Botany, Mongolian Academy of Sciences for supplying the seed samples.

References

- Arase, Y., Ikeda, K. & Murashima, N. 1997. The long-term efficiency of glycyrrhizin in chronic hepatitis C patients. *Cancer* 79: 1494-1500.
- Dashjamts, Y. 1983. *Ecological and biological characteristics of Mongolian Glycyrrhiza uralensis*. Ph.D. Dissertation. Ulaanbaatar. (in Russian)
- Gankhuyag, G. & Ligaa, U. 1993. *Mongolian Licorice*. Bayankhongor. (in Mongolian)
- Kohjyouma, M., Kohda, H., Tani, N., Ashida, K., Sugino, M., Yamamoto, A. & Horikoshi, T. 1995. *In-vitro* propagation from axillary buds of *Glycyrrhiza glabra* L. *Plant Tissue Culture Letters* 12(2): 145-149.
- Kovalenko, P.G. & Kurchii, B.A. 1998. Use of abscisic acid in the plant tissue culture of licorice *Glycyrrhiza glabra* L. electroporated protoplasts. In: *Proceedings of II International Symposium on Plant Biotechnology*, Ukraine. p. 65.
- Liao, Zh., Chen, M., Tan, F., Sun, X. & Tang, K. 2004. Micropropagation of endangered Chinese aloe. *Plant Cell Tissue and Organ Culture* 76: 83-86.
- Numazaki, K. 2003. Glycyrrhizin therapy for viral infections. *African Journal of Biotechnology* 2(10): 392-393.
- Oyunbileg, Yu. & Mijidsuren, A. 2004. Studies on tissue culture of *Glycyrrhiza uralensis* Fisch. In: *Proceedings of the Second Workshop on Korea-Mongolia Science and Technology Cooperation, Biotechnology*, Seoul, Korea, p.106-112.
- Shiraki, K., Sato, H., Kurokawa, M. & Yoshida, Y. 2003. Glycyrrhizin: the molecule and present knowledge of action in various liver diseases. In: *Prevention of progression in chronic liver disease*. (Ed. by N. Hayashi & M.P. Manns). Falk workshop, Freiburg, Germany. p.76-89.
- Thengane, S.R., Kulkarni, D.K. & Krishnamurthy, K.V. 1998. Micropropagation of licorice (*Glycyrrhiza glabra* L.) through shoot tip and nodal cultures. *In Vitro Cellular and Developmental Biology of Plants* 34(4): 331-334.

Хураангуй

Урал чихэр өвс (*Glycyrrhiza uralensis* Fisch.) нь флавоноид болон сапонины төрлийн нэгдлүүдээр баялаг, чухал ач холбогдолтой эмийн ургамлын нэг юм. Урал чихэр өвсний эдийн өсгөврийг хийхэд *in vitro*-д үрийг соёолуулсан пухуйцаас үрийн тал, үндэсний эксплант, бичил үржүүлэг хийх зорилгоор ариутгасан үе зайдмыг хэрэглэв. Чихэр өвсний каллус, нахиаг хөгжүүлэхэд ялгаатай гормоны хувилбартай Гамбург В5, Мурашиге-Скугийн хагас найрлагатай тэжээлийн орчин хэрэглэсэн юм. Лабораторийн нөхцөлд зарим бичил ургамаланцар, эсийн биомасс гаргаж авах боломжтой болох нь бидний судалгааны зарим үр дүнгээс харагдаж байна.

Received: 11 March 2005

Accepted: 16 September 2005